

Extracting CKM γ phase from $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ and

$$B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$$

Ignacio Bediaga, Gabriel Guerrer and Jussara M. de Miranda

bediaga@cbpf.br, guerrer@cbpf.br and jussara@cbpf.br

*Centro Brasileiro de Pesquisas Físicas,
Rua Xavier Sigaud 150,
22290-180 – Rio de Janeiro, RJ, Brazil*

Abstract

We discuss some aspects of the search for CP asymmetry in the three body B decays, revealed through the interference among neighbor resonances in the Dalitz plot. We propose a competitive method to extract the CKM γ angle combining Dalitz plot amplitude analysis of $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ and untagged $B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$. The method also obtains the ratio and phase difference between the *tree* and *penguin* contributions from B^0 and $\bar{B}^0 \rightarrow K^{*\pm} \pi^\mp$ decays and the CP asymmetry between B^0 and $\bar{B}^0$. From Monte Carlo studies of 100K events for the neutral mesons, we show the possibility of measuring γ .

PACS numbers:

I. INTRODUCTION

According to the Standard Model, CP violation is only possible through the CKM complex parameters. Interference between processes with different weak phases contributing to a same final state can generate an asymmetry in charge conjugate B meson decays. The most established methods to extract the CKM phases exploits the interference between the phases generated in the $B^0 \bar{B}^0$ oscillation and the phase of the decay of these particles in the same final state. Using such approaches on the $B^0 \rightarrow J/\Psi K_s$ decay, the CKM β angle is measured to a good precision. However, the other two CKM angles, α and γ , retain sizable uncertainties, due to both experimental and theoretical difficulties. LHC should determine γ to $\sim 5^\circ$ in one year of data taking[1].

Any asymmetry observed in charge conjugated B decays is interpreted as a manifestation of the CKM phases. In the two body decay scenario, direct CP asymmetry ($\mathcal{A}_{CP}$) was established by the experiments BaBar [2] and Belle [3] by simply counting the difference in number of events of the $B^0 \rightarrow K^+\pi^-$ and $\bar{B}^0 \rightarrow K^-\pi^+$ decays. In three body B meson decays, BaBar and Belle [4, 5] claim the observation of direct $\mathcal{A}_{CP}$ in the channel $B^\pm \rightarrow K^\pm \rho^0$ measured through the amplitude analysis of the final state $B^\pm \rightarrow K^\pm \pi^+ \pi^-$. They see a difference in the decay fractions among the intermediate states in the B^+ and B^- samples. That is the difference between the square of the fitted amplitude for the intermediate state integrated on the phase space. This quantity is a straightforward application of the two body strategy into the three body decays, including the requirement of a strong phase difference to observe $\mathcal{A}_{CP}$.

Beyond the fraction asymmetry quoted by Belle and BaBar, one could explore the asymmetry associated with the interference among two intermediate states, i.e. the interference between two neighbor resonances decaying to the same three body final state. As will be discussed in detail in section III, whenever the phase difference between two resonant amplitudes involves the weak phase, one should expect to see $\mathcal{A}_{CP}$ in the interference terms of the Dalitz plot distribution. This contribution to the asymmetry depends only on the existence of a weak phase difference and it can be observed even in situations where there is no strong phase difference. There are a lot of opportunities for investigation of the asymmetry in the Dalitz plot distribution rather than difference in the decay fractions alone[6, 7].

The idea of extracting γ exploring the Dalitz interference among intermediate amplitudes

in three body B decays was initially proposed for $B^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ [8], where χ_{c0} plays a fundamental role as reference channel. However, the method is statistically limited by the low contribution of the Cabibbo suppressed amplitude $B^\pm \rightarrow \chi_{c0} \pi^\pm$ [9], and also estimated through the already observed Cabibbo allowed $B^\pm \rightarrow \chi_{c0} K^\pm$ channel [4, 5]. Recently, a new effort has been made to study γ in $B \rightarrow K \pi \pi$ decays, however this approach requires time-dependent analysis [10, 11], having to deal with tagging inefficiencies that could be as large as 90% at LHCb [12].

We present a new possibility to measure the CKM γ angle, through the interference terms in the amplitude analysis of three body B decays. The method combines the experimental data from $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ and $B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$. From the charged B decays we have the *penguin* amplitude coefficient and phase of the channel $K^* \pi$. We then use this result in an untagged analysis of the neutral system, to extract the *tree* component strong phase and amplitude coefficient and the weak phase γ . The method is based on the ability to measure independently the relative amplitudes and phases for B^0 and $\bar{B}^0$ decays in a joint untagged sample.

II. AMPLITUDE ANALYSIS

The standard amplitude analysis, extensively used in non-leptonic three body heavy quark meson decays, is based on the Isobaric model. In these analyses, the two body resonant intermediate state amplitudes are represented by Breit-Wigner functions multiplied by angular distributions associated with the spin of the resonance. The various contributions are combined in a coherent sum with complex coefficients that are extracted from maximum likelihood fits to the data. The absolute value of the coefficients, normally referred to as magnitudes, are related to the relative fraction of each intermediate state and the phase takes into account rescattering or the final state interaction (FSI) between the resonance and the bachelor particle. This phase is considered constant because it depends only on the total energy of the system, i.e. the heavy meson mass. The presence of strong final state constant phases can be directly revealed through the interference between two different amplitudes. Thus, the total Isobaric amplitude, written as functions of the Dalitz plot variables, s_{12} and s_{13} is:

$$\mathcal{A}_T = a_{NR} e^{i\delta_{NR}} \mathcal{A}_{NR}(s_{12}, s_{13}) + \sum_{n=1}^N a_n e^{i\delta_j} \mathcal{A}_n(s_{12}, s_{13}). \quad (1)$$

The fit parameters are the coefficient magnitudes, a , and the phases, δ . The non-resonant amplitude, $\mathcal{A}_{NR}$ is usually represented by a constant. Each resonant amplitude, $\mathcal{A}_n (n \geq 1)$, is written as a product of three terms:

$$\mathcal{A}_n = {}^J F_n \times {}^J \mathcal{M}_n \times BW_n \quad (2)$$

The first term are the Blatt-Weisskopf damping factors [13]. ${}^J \mathcal{M}_n$ is a term which accounts for angular-momentum conservation and depends on the spin, J , of n^{th} the resonance, such that it is 1 for a spin-0, $-2 |\mathbf{p}_3| |\mathbf{p}_2| \cos\theta$ for spin-1, and $\frac{4}{3}(|\mathbf{p}_3| |\mathbf{p}_2|)^2 (3\cos^2\theta - 1)$ for spin-2, in the case of the resonance formed by the pair of particles **1** and **3**. $\mathbf{p}_2$ and $\mathbf{p}_3$ are the the 3-momentum of particles **2** and **3**, and θ is the angle between particles **2** and **3**, all measured in the resonance rest frame. The last term is a relativistic Breit-Wigner function given by:

$$BW_n = \frac{m_n \Gamma_n}{m_n^2 - s_{ij} - i m_n \Gamma_n(s_{ij})}, \quad (3)$$

where m_n is the resonance mass and $\Gamma_n(s_{ij})$ is the mass dependent width.

The relative contribution of each intermediate state is given by fractions defined as:

$$f_n \equiv \frac{\int ds_{12} ds_{13} |c_n \mathcal{A}_n|^2}{\int ds_{12} ds_{13} \sum_{jk} |c_j \mathcal{A}_j c_k^* \mathcal{A}_k^*|}. \quad (4)$$

where $\int ds_{12} ds_{13} \sum_{jk} |c_j \mathcal{A}_j c_k^* \mathcal{A}_k^*|$ is related to the total number of events of the non-leptonic three body heavy meson decay.

III. THE DIRECT DALITZ PLOT ASYMMETRY $\mathcal{A}_{DP}$

One important feature of the Dalitz plot associated with the interference between two neighboring resonances is that there will be an $\mathcal{A}_{CP}$ manifestation, independent of the strong phase differences. To illustrate this point, let us develop a simple example of two resonant scalar amplitudes, interfering in a particular region at the three body decay phase space. Suppose one of the resonant amplitudes, $\mathcal{A}_1$, has a *tree* component, with magnitude a_T and strong phase δ_T and weak phase γ , plus a *penguin* contribution with a_P and δ_P and no weak

phase, interfering with another resonant amplitude, $\mathcal{A}_R$, with magnitude and phase a_R and δ_R and the absence of weak phase. The charge conjugate total decay amplitudes are:

$$A^+ = [a_T e^{i(\delta_T + \gamma)} + a_P e^{i\delta_P}] \mathcal{A}_1 + a_R e^{i\delta_R} \mathcal{A}_R \quad (5)$$

$$A^- = [a_T e^{i(\delta_T - \gamma)} + a_P e^{i\delta_P}] \mathcal{A}_1 + a_R e^{i\delta_R} \mathcal{A}_R \quad (6)$$

and

$$\begin{aligned} |A^+|^2 - |A^-|^2 = & -4a_T a_P \sin \gamma \sin(\delta_T - \delta_P) |\mathcal{A}_1|^2 \\ & -4a_R a_T \sin \gamma (\sin(\delta_T - \delta_R) \operatorname{Re}(\mathcal{A}_1 \mathcal{A}_2^*) + \cos(\delta_T - \delta_R) \operatorname{Im}(\mathcal{A}_1 \mathcal{A}_2^*)) \end{aligned} \quad (7)$$

From Equation 7, we see that the first term has a similar behavior as the two body decays, i.e. $\mathcal{A}_{CP}$ depends on the magnitude of the *tree* component and also on the phase difference $(\delta_T^1 - \delta_P^1)$, which must be non zero for $\mathcal{A}_{CP}$ to be observable. This is the term responsible for the $\mathcal{A}_{CP}$ quoted through the difference in integrated fractions made by Babar and Belle [2] [3]. However the presence of second or third terms, the interference ones, depends only on the magnitude of the *tree* component, since the phase difference has a sine form associated with the real part of the BW's product, but also a cosine component associated with the imaginary part of the BW's product. So the interference will always produce $\mathcal{A}_{CP}$ provided that the magnitude associated with the weak phase is significant.

Interferences in the Dalitz plot do not allow to associate directly the fraction, defined in Eq. 4, with the number of events in a intermediate state channel. The relative phase between two amplitudes sharing the same phase space region can produce destructive or constructive interferences, decreasing or increasing the number of events in the three body final sample, in which case we lose deterministic information of the source of the events in the region. Due this difficulty to the concept of the direct $\mathcal{A}_{CP}$ to the three body decays, the asymmetry observed from fractions would be better referred as *quasi-direct* $\mathcal{A}_{CP}$.

The direct $\mathcal{A}_{CP}$ in the three body decays, in the sense of event counting, would be more conveniently defined by the notion of two asymmetries, one local and one global. A global $\mathcal{A}_{CP}$ is the difference between the total number of events of two conjugate three body decays. The local would be the difference in the number of events depending on the Dalitz variables s_{ij} . In general, the global asymmetry loses part of dynamics information, since it

is the integral over the phase space. On the other hand, local asymmetry accounts for the difference of the magnitude square, or the first term of Equation 7, and also the asymmetry coming from the interference terms among two resonances, or the last terms of Equation 7.

The local CP asymmetry of the charged B meson three body decay can be defined through the subtraction of the two Dalitz surfaces relative to the charged conjugated B signals:

$$\mathcal{A}_{DP} = \frac{N^+(s_{12}, s_{13}) - N^-(s_{12}, s_{13})}{N^+(s_{12}, s_{13}) + N^-(s_{12}, s_{13})} \quad (8)$$

we call $\mathcal{A}_{DP}$ the local asymmetry at the Dalitz plot, with $N^\pm(s_{12}, s_{13})$ being the number of events in s_{12}, s_{13} .

For the neutral systems with B^0 and $\bar{B}^0$ decaying to the same final state, $K_S\pi^+\pi^-$, $K_S p\bar{p}$, $\pi^0\pi^+\pi^-$ for example, Burdman and Donoghue [6] propose a convenient definition of local $\mathcal{A}_{DP}$. They showed that for a decay with $|p/q| = 1$, the asymmetry is time independent and can be written as:

$$\mathcal{A}_{dp} = \frac{N^0(s_{12}, s_{13}) - N^0(s_{13}, s_{12})}{N^0(s_{12}, s_{13}) + N^0(s_{13}, s_{12})} \quad (9)$$

A measure of local $\mathcal{A}_{DP}$ is obtained by comparing the number of events in points of the Dalitz plot symmetrical with respect to the diagonal line $s_{12} = s_{13}$.

To quantify the total amount of asymmetry present in a three body decay, one can compute the sum of the modulus of $\mathcal{A}_{DP}$, taking in account properly statistical fluctuations.

IV. METHOD

We now introduce a new possibility to measure γ , based on a time-independent analysis of $B \rightarrow K\pi\pi$ decays. The most relevant observed intermediate states for $B^\pm \rightarrow K^\pm\pi^+\pi^-$ and $B^0, \bar{B}^0 \rightarrow K_S\pi^+\pi^-$ [4, 5, 14], are summarised in Table I with the respective *tree* and *penguin* contributions, and CKM weak phases contained in the amplitude.

Based on SU(2) flavor symmetry, we expect the same *penguin* amplitudes for the four processes $B^\pm \rightarrow K^*\pi^\pm$ and $B^0, \bar{B}^0 \rightarrow K^{*\pm}\pi^\mp$. This property allows to extract the $B^\pm \rightarrow K^*\pi^\pm$ *penguin* parameters [23], and use them in the B^0 and $\bar{B}^0$ joint fit to measure the *tree* phases, from which γ is obtained.

Since we measure only relative magnitudes and phases we must elect one resonance to have fixed parameters. The compatibility between *penguin* parameters from $B^\pm$ and $B^0, \bar{B}^0$

	resonance	contribution	weak phase
$B^+ \rightarrow$	$K^*(890)^0 \pi^+$	$V_{bt} V_{ts}^* P$	
	$K^*(1430)^0 \pi^+$		
	$K^+ \rho(770)^0$	$V_{bt} V_{ts}^* P + V_{bu} V_{us}^* T_S^C$	γ
	$K^+ f_0(980)$		
	$K^+ \chi_{c0}$	$V_{bc} V_{cs}^* T_S$	
$B^0 \rightarrow$	$K^*(890)^+ \pi^-$	$V_{bt} V_{ts}^* P + V_{bu} V_{us}^* T^C$	γ
	$K^*(1430)^+ \pi^-$		
	$K_S \rho(770)^0$	$V_{bt} V_{ts}^* P + V_{bu} V_{us}^* T_S$	γ
	$K_S f_0(980)$		
	$K_S \chi_{c0}$	$V_{bc} V_{cs}^* T_S$	

TABLE I: Resonances and dominant contributions from $B \rightarrow K\pi\pi$ decays . We denote allowed, suppressed by color and *penguin* amplitudes by T^C , T_S and P .

to $K^*\pi$, requires that the amplitude analysis be made relative to an anchor resonance that has the same amplitude for B charged and neutral. For that we chose the non CP violating $B \rightarrow K \chi_{c0}$ amplitude. The asymmetry measured by Belle [15] for the channel $B^\pm \rightarrow K^\pm \chi_{c1}$ is $A_{CP} = -0.01 \pm 0.03 \pm 0.02$, indicating that the dominant contribution is a *tree* diagram without weak phase.

Following is a schematic representation of the method, with the arrows pointing its flow:

$$\begin{aligned}
B^\pm \rightarrow K^{*0} \pi^\pm & : \quad \underline{a}_P e^{i\delta_P} \\
& \Downarrow \\
B^0 \rightarrow K^{*+} \pi^- & : \quad a_P e^{i\delta_P} + \underline{a}_T e^{i\theta^+} \\
\bar{B}^0 \rightarrow K^{*-} \pi^+ & : \quad a_P e^{i\delta_P} + \underline{a}_T e^{i\theta^-} \\
& \Downarrow \\
& \gamma = \frac{\theta^+ - \theta^-}{2}.
\end{aligned} \tag{10}$$

where $\theta^\pm = (\delta_T \pm \gamma)$. We underlined the parameters measured in each amplitude analysis.

The method is based on three basic and well accepted hypothesis that can be tested during the analysis. *Hypothesis 1:* the dominant contribution of $B^\pm \rightarrow K^* \pi^\pm$ is $V_{bt} V_{ts}^* P$. This has been indicated by the BaBar result [4]. The test is to check if the $B^+ \rightarrow K^* \pi^+$ and $B^- \rightarrow K^* \pi^-$ amplitudes are the same. *Hypothesis 2:* the *penguin* components from $B^\pm \rightarrow K^* \pi^\pm$ and $B^0, \bar{B}^0 \rightarrow K^{*\pm} \pi^\mp$ are equal [16, 17]. *Hypothesis 3:* χ_{c0} have the same amplitude for $B^\pm \rightarrow K^\pm \chi_{c0}$ and $B^0, \bar{B}^0 \rightarrow K_S \chi_{c0}$. The experimental test for the second and third hypothesis, consists in the equality of the *tree* magnitudes measured from the intermediate process $B^0, \bar{B}^0 \rightarrow K^{*\pm} \pi^\mp$. The confidence in the result of γ is based in the fulfillment of all hypotheses, whereas the failure of either implies interesting unexpected effects, like a big W annihilation contribution or even a new contribution for the $B^\pm \rightarrow K^\pm \chi_{c0}$ other than the tree component.

V. $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ AND $B^0, \bar{B}^0 \rightarrow K_S \pi^+ \pi^-$ AMPLITUDES

To obtain the $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ parameters, we apply the usual maximum likelihood fit to the isobaric amplitudes given by Equation 2 with the intermediate states $i = K^*(890)^0, K^*(1430)^0, \rho(770)^0, f_0(980)$ and $K^+ \chi_{c0}$.

Regarding the neutral system, separation between B^0 and $\bar{B}^0$ to $K_S \pi^+ \pi^-$ is not trivial. The samples could be distinguished at the production level only through partner b particle tagging techniques. Furthermore, there is mixing among B^0 and $\bar{B}^0$, introducing time dependence in the decay probabilities.

The probability distribution for the final state $K_S \pi^+ \pi^-$, independent of its origin, is given by the incoherent sum: $|M(\Delta t)|^2 + |\bar{M}(\Delta t)|^2$. The matrix element for observing the original B^0 and $\bar{B}^0$ decay to the common final state are respectively:

$$M(\Delta t) = e^{-(\Gamma/2 - iM)\Delta t} [\mathcal{A} \cos(\Delta m \Delta t/2) - i q/p \bar{\mathcal{A}} \sin(\Delta m \Delta t/2)], \quad (11)$$

$$\bar{M}(\Delta t) = e^{-(\Gamma/2 - iM)\Delta t} [\bar{\mathcal{A}} \cos(\Delta m \Delta t/2) - i p/q \mathcal{A} \sin(\Delta m \Delta t/2)], \quad (12)$$

where $\mathcal{A}$ e $\bar{\mathcal{A}}$ are time-independent decay amplitudes equivalent to Equation 1, but for $B^0 \rightarrow K_S \pi^+ \pi^-$ and $\bar{B}^0 \rightarrow K_S \pi^+ \pi^-$ decays.

Assuming the same production rate for B^0 and $\bar{B}^0$, and in the case of $|p/q| = 1$, it was shown [6, 18] that this sum displays the interesting property of canceling mixing dependence

terms,

$$|M(\Delta t)|^2 + |\bar{M}(\Delta t)|^2 = e^{-\Gamma t} (|\mathcal{A}^0|^2 + |\bar{\mathcal{A}}^0|^2). \quad (13)$$

Similarly to Equation 1, $\mathcal{A}^0$ and $\bar{\mathcal{A}}^0$ are written as:

$$\mathcal{A}^0 = a_\chi e^{i\delta_\chi} \mathcal{A}_\chi + \sum_i a_i e^{i\delta_i} \mathcal{A}_i \quad (14)$$

$$\bar{\mathcal{A}}^0 = \bar{a}_\chi e^{i\bar{\delta}_\chi} \mathcal{A}_\chi + \sum_i \bar{a}_i e^{i\bar{\delta}_i} \mathcal{A}_i \quad (15)$$

As a fundamental step to achieve a γ measurement, we propose a method of *joint fitting* the set of untagged B^0 plus $\bar{B}^0$ time integrated event sample by doing a maximum likelihood fit to the Probability Distribution Function (PDF) given by

$$PDF = \frac{|\mathcal{A}^0|^2 + |\bar{\mathcal{A}}^0|^2}{N^0 + \bar{N}^0}, \quad (16)$$

where $N^0 = \int |\mathcal{A}^0|^2 ds_{ij} ds_{jk}$, and $\bar{N}^0 = \int |\bar{\mathcal{A}}^0|^2 ds_{ij} ds_{jk}$ and a, δ and $\bar{a}, \bar{\delta}$ are free fit parameters.

In the standard single sample Dalitz plot amplitude analysis, described in section II, two parameters, one magnitude and one phase, should be fixed. What is measured are the relative contributions and phase differences. The magnitude sets the overall scale. This would be the case for each of the two charged modes with a total of four fixed parameters. In the joint fit, we have to fix three parameters, one magnitude, that sets the overall scale for both modes, and two phases, one for each c.c mode. In the joint fit, the freedom of all magnitudes of one of the modes, say $\bar{B}^0$, is what guaranties that we are able to have different total number of events for each mode. In our particular case we fixed a_χ, δ_χ and $\bar{\delta}_\chi$. We are able to measure a difference in number of events from B^0 and $\bar{B}^0$, searching for direct global asymmetrie. With one single procedure it is possible to extract in an independent way the parameters from the amplitudes $\mathcal{A}^0$ and $\bar{\mathcal{A}}^0$ and the ratio in number of events, calculated by the ratio of the normalizations $N^0/\bar{N}^0$.

In the convention $(K_S, \pi^+, \pi^-) \rightarrow (\mathbf{1}, \mathbf{2}, \mathbf{3})$ for the final state particle numbering, the charge conjugation operation is equivalent to switching the Dalitz variables $s_{12} \leftrightarrow s_{13}$. The resonances K^* bands are aligned in different axis for $\mathcal{A}^0$ and $\bar{\mathcal{A}}^0$, establishing some sort of signature for the event origin B^0 or $\bar{B}^0$, as can be seen schematically in Fig. 1. Although the $\pi\pi$ resonances for B^0 and $\bar{B}^0$ overlap, the interference regions between $K\pi$ and $\pi\pi$

resonances are well separated. These interference regions are vital for establishing the phases, and they guarantee that the fitting procedure finds a unique solution. Although we cannot distinguish events, the joint fit can identify two different overlapping surfaces relative to B^0 and $\bar{B}^0$.




FIG. 1: Schematic representation for the Dalitz plot of a) $B^0 \rightarrow K_S \pi^+ \pi^-$, b) $\bar{B}^0 \rightarrow K_S \pi^+ \pi^-$ and c) the joint figure obtained for the untagged sample. Bands pictorially represent resonances regions. The vertical and horizontal bands are $K\pi$ resonances the in the diagonal are $\pi\pi$ overlapping resonances. For simplicity we ignore angular effects in the distribution. Circles are displaying regions of interference within each B^0 or $\bar{B}^0$ decay. There is no interference between the horizontal and the vertical bands for they do not belong to the same mode. It is the non-overlapping of circles that assures the separation between both amplitudes in the joint fit analysis.

VI. FEASIBILITY STUDY

To investigate the feasibility and the error size in the joint analysis, we generated and fitted one hundred Monte Carlo samples of 100K B^0 plus $\bar{B}^0$ events, that can be obtained by LHCb experiment. We do not consider any theoretical or experimental systematic uncertainties. Detector and background effects are as well not taken into account.

The parameters we used to generate the Monte Carlo samples are listed in Table II, column input. They are close to the values obtained in measured parameters for the $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ decays [4, 5]. For the B^0 and $\bar{B}^0 \rightarrow K_S \pi^+ \pi^-$ decays we used the same parameters as in the charged mode, except for including a *tree* contribution, discussed below, for $K^*(890)$

and $K_0^*(1430)^0$ and the γ phase, taken as 69° . The Dalitz plot distribution for one generated experiment of 100K events is displayed in Fig. 2, where the small contribution of the resonance χ_{c0} can be seen.




FIG. 2: $B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$ events distribution from fast MC generated with table II parameters.

One important issue concerning the γ extraction, is the size of the ratio $r = a_T/a_P$ and the phase difference $\theta = \delta_T - \delta_P$ from the *tree* and *penguin* amplitudes of the K^* resonances. The smallest the value of r the more difficult it is to apply this procedure to measure γ . The theoretical knowledge of these quantities is model dependent. Some groups use factorization approach [17], and obtain large r and small θ . On the other hand, non-factorisable approach for pseudoscalar-pseudoscalar B decay [19], presents an opposite scenario, with small r and large θ . The joint fit, applied to the real data, will be able to define which theoretical approach is more adequate, since it is possible to measure the parameters under discussion. For our study, we chose $r = 0.45$.

As stated in section IV, the first step of the method would be to solve the charged mode Dalitz Plot, from which hypothesis 1 could be tested. The parameters a_P and δ_P would then be input to the neutral system fit. Since charged mode analysis presents no novelty and the statistical errors are expected to be low compared to the neutral decays we skip this step

decay	$B^0/\bar{B}^0$	input	fit 100K events
$\chi_c K_s$	$a_0/\bar{a}_0$	0.30/0.30	fixed/(0.30 ± 0.03)
	$\delta_0/\bar{\delta}_0$	3.78/3.78	fixed/fixed
$K^*(890)\pi$	$a_1/\bar{a}_1$	1.17/1.30	1.17 ± 0.06 / (1.30 ± 0.01)
	$\delta_1/\bar{\delta}_1$	0.40/5.98	(0.41 ± 0.08)/(5.99 ± 0.07)
$K_0^*(1430)\pi$	$a_2/\bar{a}_2$	2.45/2.72	(2.45 ± 0.11)/(2.72 ± 0.13)
	$\delta_2/\bar{\delta}_2$	0.375/6.00	(0.39 ± 0.08)/(6.00 ± 0.06)
$\rho^0 K_s$	$a_3/\bar{a}_3$	0.60/0.60	(0.60 ± 0.04)/(0.60 ± 0.04)
	$\delta_3/\bar{\delta}_3$	1.20/1.20	(1.22 ± 0.09)/(1.20 ± 0.07)
$f_0 K_s$	$a_4/\bar{a}_4$	1.03/1.03	(1.02 ± 0.06)/(1.04 ± 0.05)
	$\delta_4/\bar{\delta}_4$	2.30/2.30	(2.30 ± 0.07)/(2.30 ± 0.08)
$N^0(B^0 \rightarrow K_s \pi^+ \pi^-)/N^0(\bar{B}^0 \rightarrow K_s \pi^+ \pi^-) = 0.84 \pm 0.12$			

TABLE II: Monte Carlo simulation for $B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$ decay. We generated sample with the the parameters a_i and δ_i for B^0 and $\bar{a}_i$ and $\bar{\delta}_i$ for $\bar{B}^0$. The third column display the fit results with one hundred samples for each 100K events. The last line is the ratio between $B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$ number of events.

for this study.

Each 100K event sample of B^0 plus $\bar{B}^0$ generated with the amplitudes and phases listed in the column input of Table II were fitted with an unconstrained model with just three parameters fixed. In the last column of Table II we display the central values of the Gaussian distributions of the 100 experiment's fit result and the errors are the width of the same Gaussian distribution. The total number of events of each B^0 and $\bar{B}^0$ is the result of the integration of either amplitude within the Dalitz plot. The fact that the χ_{c0} magnitude for $\bar{B}^0$ is measured to be equal to the fixed a_0 is the test for hypothesis 3. The extracted quantities are in agreement with the generated ones and with small errors, stating the feasibility of the joint fit method. Using the parameters from Table II and the relations developed in section IV, we measure $\gamma = 69^\circ \pm 7^\circ$.

If in the unconstrained analysis, the three hypotheses prove to be correct, we can reduce the numbers of free parameters assuming that the K^* resonances parameters from B^0 and

$\bar{B}^0$ are given by the following *tree* and *penguin* sum:

$$\begin{aligned} a e^{i\delta} &= a_P e^{i\delta_P} + a_T e^{i\delta_T + \gamma}, \\ \bar{a} e^{i\bar{\delta}} &= a_P e^{i\delta_P} + a_T e^{i\delta_T - \gamma}. \end{aligned} \quad (17)$$

Instead of fitting for $a, \delta, \bar{a}$ and $\bar{\delta}$, we fit for a_T, δ_T and γ since we can fix a_P, δ_P from $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ result. This way, constraining the fit with the hypothesis previously tested we reduce the error in γ . We measure with the constrained hypothesis, $\gamma = 69^\circ \pm 5^\circ$ and also we obtained for the ratio between the *tree* and *penguin* component $r = 0.45 \pm 0.05$.

VII. CONCLUSION

We discussed some aspects of the search of $\mathcal{A}_{CP}$ in three body B decays, emphasising the main characteristic of these decays, which is the interference among two neighbor resonances in the Dalitz plot. In particular, we discuss how the CKM γ phase can be extracted applying the Isobar model[20], extensively used in many charm meson three body decay amplitude analysis. We showed that the phase differences, coming from the amplitude analysis, can provide richer information than treating the three body decays as a straightforward extension of two body hadronic decays. We showed that if there is a weak phase difference between two intermediate state, $\mathcal{A}_{CP}$ will be present in the Dalitz plot regardless of the strong phase difference.

We presented a method to extract the CKM γ angle using a combined Dalitz plot analysis from $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ and $B^0, \bar{B}^0 \rightarrow K_s \pi^+ \pi^-$. This method use three basic hypothesis that can be tested before one proceeds to the constrained fit. For measuring the B neutral analysis, we use a new technique of joint fit that allows us to extract, in an independent way, the amplitudes from two summed surfaces in a joint sample of $B^0 + \bar{B}^0$ untagged events. We carried a simplified fast MC study to estimate the errors associated to the joint fit technique. Assuming a relatively low statistics for the anchor resonance χ_{c0} we obtained γ with a 5° error.

During the analysis we can measure CP violation by counting the number of events from B^0 and $\bar{B}^0$, information extracted in the joint fit, or exploring Dalitz symmetries as discussed in [6]. We can also measure the ratio and phase difference from *tree* and *penguin* amplitudes of the K^* resonance, defining which theoretical approach, factorisable or non-factorisable,

is more adequate.

The method presented here is competitive with the other approaches to determine the CKM γ angle [1] and needs the high statistics expected for the LHCb experiment, due to the small contribution of the reference channel $B \rightarrow \chi^0 K$. However, in the case that the $f_0(980)$ resonance is dominated by the $s\bar{s}$ component [21], or even if the ratio between the tree and *penguin* is negligible [22], the $B \rightarrow f_0(980)K$ amplitude could take the place of the charmonium as a reference channel in the analysis.

Acknowledgements

We thank Professor Giovanni Bonvicini for suggestions and comments.

-
- [1] O. Schneider, talk at the " Flavor in the era of the LHC" workshop, CERN, November 2005, <http://cern.ch/flavlhc> and, J. Rademacker, talk at the "Physics at the LHC" workshop, Krakow, July 2006, <http://indico.cern.ch/conferenceDisplay.py?confId=a058062>.
 - [2] BaBar Collaboration, B. Aubert et al., Phys. Rev. Lett. **93**, 131801 (2004).
 - [3] Belle Collaboration, Y. Chao et al., Phys. Rev. Lett. **93**, 191802 (2004).
 - [4] BaBar Collaboration, B. Aubert *et al.*, Phys.Rev.**D72**:072003, (2005), Erratum-*ibid.***D74**:099903, (2006). and hep-ex/0507004.
 - [5] Belle Collaboration, A. Garmash *et al.*, Phys. Rev. Lett. **96** 251803 (2006), hep-ex/0512066.
 - [6] G. Burdman and J.F. Donoghue, Phys. Rev. **45** 187 (1992)
 - [7] CLEO Collaboration, D. Asner *et al.*, Phys.Rev. **D70**, 091101 (2004).
 - [8] I. Bediaga, R. E. Blanco, C. Göbel and R. Méndez-Galain, Phys. Rev. Lett. **81** 4067 (1998).
 - [9] BaBar Collaboration, B. Aubert et al., Phys.Rev. **D72**:052002, (2005), Phys.Rev. **D76**:011102,(2007) and hep-ex/0507025.
 - [10] M. Ciuchini, M. Pierini, L.Silvestrini,Phys.Rev.**D74**:051301,(2006) hep-ph/06010233.
 - [11] M. Gronau, D. Pirjol, A. Soni, J. Zupan,Phys.Rev.**D75**:014002,(2007) and hep-ph/0608243.
 - [12] LHCb Collaboration, Reoptimized detector design and performance report, CERN-LHCC/2003-030.
 - [13] J. M. Blatt and V. F.Weisskopf, Wiley, New York p. 361 (1952)
 - [14] A. Garmash, representing Belle Collaboration. Proceeding of 11⁰ International Conference

- on Hadron Spectroscopy, Rio de Janeiro August of 2005. AIP Conference Proceeding and hep-ex/0510059. Also in K. Abe *et al.*, Phys.Rev. **D75**, 012006 (2007) and hep-ex/0610081.
- [15] Belle Collaboration, R. Kumar *et al.*, Phys.Rev.**D74**:051103, (2006) and hep-ex/0607008.
- [16] C. W. Chiang *et. al.*, Phys. Rev. **D69**, 034001 (2004).
- [17] M. Beneke, M. Neubert, Nucl. Phys., **B675**, 333 (2003).
- [18] S. Gardner, J.Tandean, Phys.Rev. **D69**, 034011 (2004).
- [19] A. J. Buras *et al.*, Phys.Rev.Lett.**92**:101804 (2004).
- [20] Yao, W.-M. J. of Physics G **33**, 1 (2006). In particulary the D. Asner review *Dalitz plot analysis formalism* pag. 713
- [21] I. Bediaga, F.S. Navarra and M. Nielsen Phys. Lett. **579 B**, 59 (2004).
- [22] H. Cheng and K. Yang, Phys.Rev.**D71**:054020, 2005 and A.K. Giri, B. Mawlong and R. Mohanta, hep-ph/0608088.
- [23] For simplicity we refer only to K^* , while in practical the method uses simultaneously the parameters of the resonances $K^*(890)$ and $K^*(1430)$.